

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 687 411 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 95109088.5

(51) Int. Cl.<sup>6</sup>: **A01N 1/02**

(22) Date of filing: 13.06.95

(30) Priority: 14.06.94 JP 131654/94


(43) Date of publication of application:  
20.12.95 Bulletin 95/51(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC  
NL PT SE**(71) Applicant: **Senju Pharmaceutical Co., Ltd.**  
5-8, Hiranomachi 2-chome,  
Chuo-Ku  
Osaka-shi,  
Osaka 541 (JP)(72) Inventor: **Ishimura, Yuzuru**  
Karsa Mitaka 104,  
7-7-14, Shimorenjaku

**Mitaka-shi,  
Tokyo 181 (JP)**  
Inventor: **Suematsu, Makoto**  
26-4-301, Daikyocho  
Shinjuku-ku,  
Tokyo 160 (JP)  
Inventor: **Yoshida, Kenichi**  
2-16, Sojiji 2-chome  
Ibaraki-shi,  
Osaka 567 (JP)

(74) Representative: **von Kreisler, Alek,**  
**Dipl.-Chem. et al**  
**Patentanwälte,**  
**von Kreisler-Selting-Werner,**  
**Bahnhofsvorplatz 1 (Deichmannhaus)**  
**D-50667 Köln (DE)**

(54) **Hepatic graft preservative composition and a method for viable preservation of the hepatic graft**

(57) This invention provides a hepatic graft preservative composition containing a phosphoric acid diester compound of the formula:



(wherein R<sub>1</sub> and R<sub>2</sub> are the same or different and each represents a hydrogen atom or a methyl group) or a pharmacologically acceptable salt thereof and a method for viable preservation of the hepatic graft using the above substance.

The hepatic graft preservative composition and hepatic graft preserving method of this invention inhibit hepatic microcirculation disturbance following cold storage-reperfusion of the liver to effectively prevent necrosis of hepatocytes and is, therefore, useful for the preservation of the hepatic graft.

**EP 0 687 411 A1**

**BACKGROUND OF THE INVENTION****1. Field of the Invention**

5 This invention relates to an improved hepatic graft preservative composition and an improved method for viable preservation of the hepatic graft. More particularly, this invention relates to a viable liver preservative composition comprising an ascorbyl tocopheryl phosphate compound or a pharmacologically acceptable salt thereof and a method for viable preservation of the hepatic graft which comprises using said compound or salt.

**2. Description of the Prior Art**

10 For a successful organ transplantation, the organ resected from a donor must be kept functionally intact until it has been transplanted into a recipient. Each kind of organ has its own characteristics and, hence, requires a unique approach towards viable storage. However, the preservation principles applicable to all organs in common are to suppress metabolic consumption and to preserve tissue ATP levels.

For the viable preservation of the liver for transplantation, there are two methods, namely the "simple cold storage method", which consists of initial perfusion with a preservative solution at 4°C and subsequent immersion in a similar cold preservative solution and the "continuous hypothermic perfusion storage method", which involves the use of a machine. Clinically, however, the simple cold storage method using a preservative solution is generally employed.

As the liver preservative, University of Wisconsin (UW) solution is generally employed. However, this solution has the disadvantage that the reduced glutathione contained as an active ingredient is very labile and easily oxidized in aqueous medium and, therefore, must be extemporaneously added, thus complicating the procedure. Moreover, UW solution is viscous and cannot be easily buffered so that it is not fully satisfactory for preservation of the hepatic graft. Moreover, with the conventional preservative solutions including said UW solution, hepatocytes tend to be damaged for some reasons or others so that deterioration of liver function cannot be successfully prevented.

Therefore, research and development efforts are being made in earnest for developing an improved hepatic graft preservative solution free from the above-mentioned disadvantages and insuring good graft preserving effects.


15 In the course of their ceaseless research into the pharmacology of ascorbyl tocopheryl phosphate compounds, the inventors of the current application discovered that these compounds have an inhibitory effect on microcirculation disorders of the liver and are useful for the viable preservation of the liver isolated for transplantation. This discovery was followed by further studies which have resulted in the development of this invention.

**SUMMARY OF THE INVENTION**

20 This invention provides a hepatic graft preservative composition containing a phosphoric acid diester compound and a new protocol for viable preservation of the hepatic graft which involves the use of said compound.

This invention is, therefore, directed to:

25 (1) A hepatic graft preservative composition comprising a phosphoric acid diester compound of the following formula or a pharmacologically acceptable salt thereof (hereinafter referred to briefly as the compound).



- (wherein  $R_1$  and  $R_2$  are the same or different and each represents a hydrogen atom or a methyl group).
- (2) The hepatic graft preservative composition as defined under (1) as provided in a liquid dosage form.
- (3) The hepatic preservative composition as defined under (2) wherein said phosphoric diester compound or pharmacologically acceptable salt occurs in a concentration of  $5 \times 10^{-9}$  g/ml to  $1 \times 10^{-4}$  g/ml.
- (4) The hepatic graft preservative composition as defined under (2) wherein said liquid dosage form has an osmolarity of 260 mOsm to 360 mOsm.
- (5) The hepatic graft preservative composition as defined under (2) wherein said liquid dosage form has a pH value of 3 to 10.
- (6) A method of preserving the hepatic graft which comprises using a liquid composition comprising a phosphoric acid diester compound or a pharmacologically acceptable salt as defined under (1) as ① an initial perfusate for intrahepatic perfusion at resection of the liver, ② a maintenance immersion or perfusion preservative and/or ③ a rinse for use prior to warm blood reperfusion.

#### BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 is a diagrammatic representation of necrotized liver cell counts (propidium iodide (PI) fluoresceine-positive nuclei; ( $\times 10^2/\text{mm}^2$ ) in the periportal zone (zone 1), midzonal zone (zone 2) and pericentral zone (zone 3) of the rat liver after cold storage-reperfusion in the control group and the compound-treatment group.

#### DETAILED DESCRIPTION OF THE INVENTION

The term "hepatic graft preservative composition" is used in this specification to mean not only said initial perfusate but also said rinse unless contrary to the context in which the term appears.

The compound for use in the hepatic graft preservative composition and hepatic graft preserving method of this invention can be synthesized by *inter alia* the process in Japanese Patent Publication H-2-44478 or Japanese Patent Application Kokai S-62-205091, or any improvement thereof.

The compound for use in the hepatic graft preservative composition and hepatic graft preserving method of this invention is already known to be useful as an anticataract agent, a prophylactic and therapeutic agent for climacteric disturbance, a skin care cosmetic ingredient (Japanese Patent Publication H-2-44478), an anti inflammatory agent (Japanese Patent Publication H-1-27044), an antiulcer agent (Japanese Patent Application Kokai S-63-270626), a prophylactic and therapeutic agent for ischemic organic impairment (Japanese Patent Application Kokai H-2-111722) and a Maillard reaction inhibitor (Japanese Patent Application Kokai H-3-161444), among others.

The compound for use in the hepatic graft preservative composition and hepatic preserving method of this invention may be whichever of its free form and in the form of a pharmacologically acceptable salt thereof. The pharmacologically acceptable salt that can be used includes salts with alkali metals such as sodium, potassium, etc. and salts with alkaline earth metals such as calcium and magnesium, among others. Other types of salts, if acceptable from pharmacologic points of view, can also be employed.

According to the clinical objective and need, more than one species of the compound can be incorporated, in an appropriate combination, in the hepatic graft preservative composition of this invention.

The compound for use as the active ingredient in the hepatic graft preservative composition and hepatic graft preserving method of this invention is a very safe substance with an extremely low toxic potential and, as such, can be used with advantage for the purposes of this invention. [e.g. the  $\text{LD}_{50}$  values of L-ascorbyl

DL- $\alpha$ -tocopheryl phosphate potassium (hereinafter referred to briefly as EPC-K)  $\geq 5$  g/kg p.o. (rats) and  $\geq 100$  mg/kg i.v. (rats)).

The liver preservative composition of this invention can be provided in a liquid form or supplied in a solid form for extemporaneous reconstitution. The solid form can be advantageously used as dissolved, suspended or emulsified in purified water, physiological saline or like medium. The solid form includes tablets, granules and powders, among others, which can be respectively manufactured by the known techniques. These preparations may contain conventional additives such as an excipient, binder, disintegrator, dispersant, reabsorption promoter, buffer, surfactant, solubilizer, preservative, emulsifier, isotonicizing agent, stabilizer, pH control agent and so on.

Unless contrary to the object of this invention, the hepatic graft preservative composition of this invention may contain other organ-preserving ingredients which are generally used for the viable preservation of hepatic grafts. Among such ingredients are antibiotics, insulin, carbohydrates (glucose, mannitol, etc.), vitamins (vitamin C, vitamin E, etc.), organic acids (lactic acid, citric acid, etc.), nucleic acid bases (adenosine triphosphate etc.), antihypertensive agents (calcium-channel blockers,  $\beta$ -adrenergic antagonists, angiotensin-converting enzyme inhibitors, etc.), antiplatelet factor, antidiuretic hormone, anticoagulant (e.g. heparin) and so on.

Furthermore, the compound can be dissolved in any known organ preservative solution, such as Euro-Collins (EC) solution and University of Wisconsin solution [ViaSpan (registered trademark) produced by DuPont], to provide a hepatic graft preservative solution.

The proper concentration of the compound in the hepatic graft preservative composition of this invention is dependent on the species of compound, condition of the liver, and the necessary duration of preservation but the recommended final concentration in a liquid preparation is generally about  $5 \times 10^{-9}$  g/ml to  $1 \times 10^{-4}$  g/ml and preferably about  $1 \times 10^{-7}$  g/ml to  $1 \times 10^{-6}$  g/ml.

The osmolarity of such a liquid hepatic graft preservative composition of this invention is adjusted, by known means, to about 260 mOsm to about 360 mOsm, preferably about 275 mOsm to about 320 mOsm. The pH of the liquid preparation should also be adjusted, by known means, to about 3 to 10, preferably about 4 to 9.

The temperature suited for the hepatic graft preservation employing the preservative composition of this invention is dependent on the species and concentration of compound, condition of the liver and the desired duration of preservation but is generally about  $-5^{\circ}\text{C}$  to  $20^{\circ}\text{C}$ , preferably about  $0^{\circ}\text{C}$  to about  $15^{\circ}\text{C}$  and, for still better results, about  $4^{\circ}\text{C}$ .

In preserving the isolated liver with the hepatic graft preservative composition of this invention, the known organ cassette, module and other hardware can be utilized.

The hepatic graft preservative composition of this invention can be employed ① as initial perfusate for intrahepatic perfusion at resection of the liver, ② as a maintenance immersion or perfusion storage solution, and/or ③ as a rinse preceding warm blood reperfusion. Thus, using the liquid hepatic graft preservative composition of this invention within the temperature range mentioned above, the fresh liver is subjected initial perfusion from the portal vein for about 1 to 10 minutes to wash out the blood from the tissue. Then, this initially perfused graft is maintained in, or under perfusion with, a suitable amount of the liquid preservative composition of this invention within the temperature range mentioned hereinbefore. Finally, prior to reperfusion with the warm blood after transplantation, the liver is rinsed with the liquid hepatic graft preservative composition of this invention.

## EXAMPLES

The following example and formulation examples are intended to illustrate this invention in further detail and should by no means be construed as defining the scope of the invention.

### Example 1 Hepatocyte-protecting effect of the compound in the rat liver cold storage-reperfusion injury model

The hepatocyte-protecting effect of the compound in the rat liver cold storage-reperfusion injury model was studied using a rat perfused liver microcirculation monitoring system developed by the inventors of this invention.

[Test substance] L-Ascorbyl DL- $\alpha$ -tocopheryl phosphate potassium (abbreviation: EPC-K).

[Method] Livers from male Wistar rats with body weights 300 g to 330 g were perfused with Solution I<sup>1)</sup> to wash out the blood thoroughly and stored in a control group (Solution II<sup>2)</sup>) and an EPC-K group (Solution II containing  $4 \times 10^{-6}$  M EPC-K) at  $4^{\circ}\text{C}$  for 90 minutes. After this cold storage, Solution I containing  $1 \mu\text{M}$

propidium iodide (PI) was injected at a flow rate of 3.0 ml/min/g liver weight in one direction from the portal vein and after one hour of this perfusion the injury to the cells (severity of cell necrosis) in the hepatic microcirculation was assessed in terms of PI-positive nuclei, i.e. injured nuclei forming a fluorescent complex, by the technique of liver microcirculation viviperception. The stained nuclear image was input as a video picture using a high-sensitivity imaging apparatus (Silicon Intensified Target Camera, Hamamatsu Photonics) through an inverted type biomicroscope and the number of PI-positive nuclei per unit area was analyzed in the three strata of (1) periportal zone (zone 1), (2) midzonal zone (zone 2), and (3) pericentral zone (zone 3) according to the method described in the literature (Lab. Invest. 67, 434-442, 1992). Thus, one hour after initiation of reperfusion, fluorescein isothiocyanate-labelled albumin (2%, 100  $\mu$ l, bolus) was infused into the portal vein and the locations of the terminal branch of the portal vein (PV) and the central vein (CV) were determined. The area within a circle drawn with a radius of 70  $\mu$ m about PV and that about CV were defined as zone 1 and zone 3, respectively, while the remaining intermediate zone was defined as zone 2. A circle of the same radius was marked off at random within zone 2 and the number of PI-positive nuclei within the circle was counted as data for zone 2.

1) 95%O<sub>2</sub>/5%CO<sub>2</sub>-saturated Krebs-Henseleit solution

Sodium chloride	118.5 mM
Potassium chloride	4.7 mM
Calcium chloride	2.5 mM
Potassium dihydrogen phosphate	1.2 mM
Magnesium sulfate	1.2 mM
Sodium hydrogen carbonate	20.0 mM

2) Euro-Collins solution

Potassium monohydrogen phosphate	740 mg
Potassium dihydrogen phosphate	205 mg
Potassium chloride	112 mg
Sodium hydrogen carbonate	84 mg
Glucose	3.5 g
Water for injection	q.s.
	Total 100 ml

[Results] The results are shown in Fig. 1.

It is apparent from Fig. 1 that the composition of this invention inhibits the liver microcirculation disturbance after rat liver cold storage-reperfusion to effectively prevent necrosis of liver cells, suggesting that the composition is of value as a hepatic graft preservative.

## Example 2

EPC-K	0.3 mg
Potassium chloride	0.12 g
Calcium chloride	15 mg
Magnesium chloride	5 mg
Sodium chloride	0.8 g
1N-Hydrochloric acid	q.s.
1N-Sodium hydroxide	q.s.
Water for injection	q.s.
	Total 100 ml
	pH 7.3

The above ingredients are mixed in the conventional manner to provide a hepatic graft preservative solution.

**Example 3**

EPC-K	0.1 mg
Potassium dihydrogen phosphate	0.25 g
Sodium citrate	0.1 g
Potassium chloride	0.85 g
Magnesium chloride	15 mg
Calcium chloride	15 mg
Sodium chloride	0.1 g
1N-Hydrochloric acid	q.s.
1N-Sodium hydroxide	q.s.
Water for injection	q.s.
	Total 100 ml
	pH 7.3

The above ingredients are mixed in the conventional manner to provide a liver graft preservative solution.

**Example 4**

EPC-K	0.1 g
Mannitol	5 g
Water for injection	q.s.
Sodium hydroxide	q.s.
	Total 100 ml
	pH 7.3

The above ingredients are mixed in the conventional manner and filled in 2 ml-glass ampules to provide an injectable solution.

This solution is extemporaneously mixed with an appropriate amount of an organ preservative solution, such as Euro-Collins or ViaSpan solution, to provide a hepatic graft preservative solution.

**Example 5**

EPC-K	10 mg
Sucrose	500 mg

The above solid composition is filled in 5 ml-glass vials in the conventional manner to provide a hepatic graft preservative.

This preservative is extemporaneously dissolved in a suitable amount of an organ preservative solution, such as Euro-collins or ViaSpan solution, to provide a hepatic graft preservative solution.


**Example 6**

Using about 500 to 2000 ml of the hepatic graft preservative solution (4 °C ) prepared in Example 3, the fresh cadaver liver was subjected to initial perfusion for about 2 minutes and then stored in the same preservative solution at 4 °C.

The hepatic graft preservative composition and hepatic graft preserving method of this invention inhibit hepatic microcirculation disturbance following cold storage-reperfusion of the liver to effectively prevent necrosis of hepatocytes and is, therefore, useful for the preservation of the hepatic graft.

## Claims

1. A hepatic graft preservative composition comprising a phosphoric acid diester compound of the following formula or a pharmacologically acceptable salt thereof.



- (wherein  $R_1$  and  $R_2$  are the same or different and each represents a hydrogen atom or a methyl group).
- The hepatic graft preservative composition of claim 1 as provided in a liquid dosage form.
  - The hepatic preservative composition of claim 2 wherein said phosphoric diester compound or pharmacologically acceptable salt occurs in a concentration of  $5 \times 10^{-9}$  g/ml to  $1 \times 10^{-4}$  g/ml.
  - The hepatic graft preservative composition of claim 2 wherein said liquid dosage form has an osmolality of 260 mOsm to 360 mOsm.
  - The hepatic graft preservative composition of claim 2 wherein said liquid dosage form has a pH value of 3 to 10.
  - A method of preserving a hepatic graft which comprises using a liquid composition comprising a phosphoric acid diester compound or a pharmacologically acceptable salt according to claim 1 as ① an initial perfusate for intrahepatic perfusion at resection of the liver, ② a maintenance immersion or perfusion preservative and/or ③ a rinse for use prior to warm blood reperfusion.




Fig.1





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 95 10 9088

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)		
X	CHEMICAL ABSTRACTS, vol. 118, no. 1, 4 January 1993 Columbus, Ohio, US; abstract no. 4606, M.OBATA 'Investigation of lipid peroxidation in regenerating rat liver' * abstract * & NIPPON GEKA GAKKAI ZASSHI, vol. 93, no. 8, 1992 pages 833-841, ---	1-6	A01N1/02		
Y	US-A-5 306 713 (SUETSUGU KEIKO ET AL) 26 April 1994 * the whole document * ---	1-6			
Y	GB-A-2 249 937 (PASTEUR MERIEUX SERUMS VACC) 27 May 1992 * page 1, line 3 - page 2, line 18 * * page 4, line 27 - page 5, line 3 * ---	1-6			
Y	CHEMICAL ABSTRACTS, vol. 119, no. 5, 2 August 1993 Columbus, Ohio, US; abstract no. 40876, K.TANEMOTO ET AL. 'Beneficial effect of EPC-K1 on the survival of warm ischemic damaged graft in rat cardiac transplantation' * abstract * & ACTA MED OKAYAMA, vol. 47, no. 2, 1993 pages 121-127, --- -/--	1-6	<table border="1"><thead><tr><th>TECHNICAL FIELDS SEARCHED (Int.Cl.6)</th></tr></thead><tbody><tr><td>A01N</td></tr></tbody></table>	TECHNICAL FIELDS SEARCHED (Int.Cl.6)	A01N
TECHNICAL FIELDS SEARCHED (Int.Cl.6)					
A01N					
The present search report has been drawn up for all claims					
Place of search THE HAGUE		Date of completion of the search 27 September 1995	Examiner Lamers, W		
<table><tr><td><b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</td><td>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons ..... &amp; : member of the same patent family, corresponding document</td></tr></table>				<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons ..... & : member of the same patent family, corresponding document
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons ..... & : member of the same patent family, corresponding document				



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 95 10 9088

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	CHEMICAL ABSTRACTS, vol. 120, no. 7, 14 February 1994 Columbus, Ohio, US; abstract no. 74626, F.TAKAYAMA ET AL. 'Effects of anti-free radical interventions on phosphatidylcholine hydroperoxide in plasma after ischemia-reperfusion in the liver of rats' * abstract * & BIOCHEM.PHARMACOL., vol. 46, no. 10, 1993 pages 1749-1757, ---	1-6	<b>TECHNICAL FIELDS SEARCHED</b> (Int.Cl.6)
Y	EP-A-0 127 471 (SENJU PHARMA CO) 5 December 1984 * page 2, line 6 - line 15 * * page 5, line 5 - line 10 * ---	1-6	
Y	EP-A-0 339 486 (TAKEDA CHEMICAL INDUSTRIES LTD) 2 November 1989 * page 2, line 9 - line 45 * * page 3, line 33 - line 37 * * page 3, line 42 - line 43 * * page 5, line 13 * ---	1-6	
A	EP-A-0 324 387 (TAKEDA CHEMICAL INDUSTRIES LTD ; SENJU PHARMA CO (JP)) 19 July 1989 * the whole document * ---	1-6	
A	EP-A-0 590 514 (SENJU PHARMA CO) 6 April 1994 * the whole document * -----	1-6	
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>27 September 1995</b>	Examiner <b>Lamers, W</b>
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			

EPO FORM 1500 03.92 (P4/C01)